



Climate Change Assessment for the Motuoane Production Right Application (PR016), Free State Province – Scoping Phase

Project done on behalf of **Environmental Impact Management Services (Pty) Ltd**

Project Compiled by:

G Petzer

Report No: 25EIM04CS | **Date:** November 2025



Address: 62 Constantia Avenue, Mnandi AH, Centurion, 0157
Postal: PostNet Suite #18 Private Bag x59, Halfway House, 1685
Tel: +27 (0)11 805 1940

Report Details

Status	Rev 1
Report Title	Climate Change Assessment for the Motuoane Production Right Application (PR016), Free State Province – Scoping Phase
Report Number	25EIM04CS
Date	November 2025
Client	Environmental Impact Management Services (Pty) Ltd
Prepared by	Gillian Petzer, (Pr. Eng.), BEng (Chemical) (University of Pretoria)
Reviewed by	Renee von Gruenewaldt, (Pr. Sci. Nat.), MSc (University of Pretoria)
Notice	Airshed Planning Professionals (Pty) Ltd is a consulting company located in Midrand, South Africa, specialising in air quality, climate change and noise impact assessments. The company originated in 1990 as Environmental Management Services, which amalgamated with its sister company, Matrix Environmental Consultants, in 2003.
Declaration	Airshed is an independent consulting firm with no interest in the project other than to fulfil the contract between the client and the consultant for delivery of specialised services as stipulated in the terms of reference.
Copyright Warning	Unless otherwise noted, the copyright in all text and other matter (including the manner of presentation) is the exclusive property of Airshed Planning Professionals (Pty) Ltd. It is a criminal offence to reproduce and/or use, without written consent, any matter, technical procedure and/or technique contained in this document.

Revision Record

Revision Number	Date	Reason for Revision
Rev 0	August 2025	For internal review
Rev 1	November 2025	Minor edits

EXECUTIVE SUMMARY

Airshed Planning Professionals (Pty) Ltd (Airshed) was commissioned by Environmental Impact Management Services (Pty) Ltd (EIMS) to undertake a Climate Change Assessment (CCA) for the proposed Motuoane Production Right Application (PR016) (hereafter referred to as the project). This report details the scoping phase of the CCA undertaken for the project which will focus on the physical risks of climate change on the region. The climate change specialist study, which will follow the scoping phase, will assess the significance of impacts of the project greenhouse gas (GHG) emissions generated, along with the potential impact of climate change on the operation of the project.

The physical risks of climate change on the study area (based on the Fifth Assessment Report (AR5) by the Intergovernmental Panel on Climate Change (IPCC) data) can be summarised as follows:

- Climate:
 - Temperature:
 - Baseline: 3 hot days (50th percentile) and annual average temperature of 15.9°C;
 - High mitigation climate situation (RCP4.5): increase in hot days by 8.5 days with an increase in temperature of 2.3°C (50th percentile);
 - Low mitigation climate situation (RCP8.5): increase in temperature of 2.6°C (50th percentile).
 - Rainfall:
 - Baseline: average annual rainfall of 1099 mm;
 - High mitigation climate situation (RCP4.5): increase of 2.1 extreme rainfall days with an increase in rainfall of 163 mm (50th percentile);
 - Low mitigation climate situation (RCP8.5): increase of 2.2 extreme rainfall days with an increase in rainfall of 226 mm (50th percentile).
- Hazards assuming the low mitigation climate situation (RCP8.5):
 - Wildfires: Isolated pockets of moderate and high increased risk of wildfires at settlements in the municipalities, but not within the project area. The projection of 23 fire danger days per year over the project area;
 - Drought: Isolated pockets at settlements within the municipalities are at very low risk of drought with the Standardized Precipitation Index (SPI) of -0.44 for the project area;
 - Exposure to heat extremes: some settlements within the municipalities are at high risk of encountering increasing heat stresses; and,
 - Flooding: the settlements within the study area are at slight to moderate risk of increased extreme rainfall days with isolated pockets of very low to moderate increase in exposure to urban flooding for the municipalities.

TABLE OF CONTENTS

1	INTRODUCTION	1
1.1	Scope of Work	1
2	REGULATORY CONTEXT AND IMPACT ASSESSMENT CRITERIA	3
2.1	International Agreements	3
2.2	South African National Climate Change Response Policy	4
2.3	Nationally Determined Contribution	5
2.4	Greenhouse Gas Emissions Reporting	7
2.5	GHG Inventories	7
2.5.1	National GHG Emissions Inventory	7
2.5.2	GHG Emission Inventory for the Sector	8
2.5.3	Draft National Guideline for Consideration of Climate Change in Development Applications, June 2021	8
3	PHYSICAL RISKS OF CLIMATE CHANGE ON THE REGION	10
3.1	Vulnerability	10
3.2	Climate	11
3.2.1	Baseline Climate	11
3.2.2	Projected Future Climate	15
3.3	Hazards	21
3.4	Vegetation Disturbance	28
3.5	Impact of Climate Change	28
3.5.1	Water Supply	28
3.5.2	Surface Water	31
3.5.3	Ground Water	35
3.5.4	Economy	36

3.5.5	Agriculture, Forestry and Fisheries	38
3.5.6	Other Resources	40
4	IMPACT SIGNIFICANCE BASED ON SCOPING PHASE ASSESSMENT	46
5	PLAN OF STUDY FOR THE IMPACT ASSESSMENT	47
6	REFERENCES	48

LIST OF TABLES

Table 2-1: South Africa's NDC mitigation targets.....	6
Table 3-1: Forecasted economic gains or losses for the Matjhabeng Municipality for the RCP4.5 and RCP8.5 scenarios.....	38
Table 3-2: Forecasted economic gains or losses for the Masilonyana Municipality for the RCP4.5 and RCP8.5 scenarios.....	38
Table 3-3: Economic contribution of main commodities for Matjhabeng Municipality	39
Table 3-4: Economic contribution of main commodities for Masilonyana Municipality.....	39
Table 3-5: Projected economic contribution of main commodities for Matjhabeng Municipality	40
Table 3-6: Projected economic contribution of main commodities for Masilonyana Municipality	40
Table 3-7: The impacts of climate change on other resources	41

LIST OF FIGURES

Figure 1-1: Location of the project site	2
Figure 3-1: Baseline (1961 to 1990) annual average temperature and rainfall for the project area (CSIR, 2025) (dark blue marker indicates approximate location of the project)	12
Figure 3-2: Baseline (1961 to 1990) number of very hot days (>35°C) and heat wave days annually for the project area (CSIR, 2025) (dark blue marker indicates approximate location of the project)	13
Figure 3-3: Annual average temperature (top panel) and temperature anomaly (lower panel) between 1979 and 2024 (Meteoblue, 2025)	14
Figure 3-4: Annual average rainfall (top panel) and rainfall anomaly (lower panel) between 1979 and 2024 (Meteoblue, 2025)	15
Figure 3-5: Projected change in annual average temperature for the near future (2021 – 2050) (dark blue marker indicates approximate location of the project)	17
Figure 3-6: Projected change in annual average number of very hot days for the near future (2021 – 2050) (dark blue marker indicates approximate location of the project)	18
Figure 3-7: Projected change in annual average rainfall for the near future (2021 – 2050) (dark blue marker indicates approximate location of the project)	19
Figure 3-8: Projected change in annual average number of extreme rainfall days (>20 mm in <24 hours) for RCP4.5 trajectory (dark blue marker indicates approximate location of the project)	20
Figure 3-9: Risk of increased wildfires for the Matjhabeng and Masilonyana Municipalities in 2050 based on RCP8.5 trajectory (dark blue marker indicates approximate location of the project)	22
Figure 3-10: Risk of increased drought tendencies for the Matjhabeng and Masilonyana Municipalities in 2050 based on RCP8.5 trajectory (dark blue marker indicates approximate location of the project)	23
Figure 3-11: Risk of increased heat extremes for the Matjhabeng and Masilonyana Municipalities in 2050 based on RCP8.5 trajectory (dark blue marker indicates approximate location of the project)	24
Figure 3-12: Risk of increased flooding for the Matjhabeng and Masilonyana Municipalities in 2050 based on RCP8.5 trajectory (dark blue marker indicates approximate location of the project)	25
Figure 3-13: Current water stress for the project area (Hofste, et al., 2019) (blue dot indicates project location)	26
Figure 3-14: Projected (2050) water stress for the project area (Hofste, et al., 2019) (blue dot indicates project location)	27
Figure 3-15: Current water availability for the Matjhabeng Municipality	29

Figure 3-16: Current water availability for the Masilonyana Municipality	29
Figure 3-17: Estimated current and future (2050) water supply vulnerability based on medium population growth for the Matjhabeng Municipal	30
Figure 3-18: Estimated current and future (2050) water supply vulnerability based on medium population growth for the Masilonyana Municipal	31
Figure 3-19: Quaternary catchment areas for the study area	32
Figure 3-20: Current annual and monthly surface water runoff, precipitation and evaporation for the Matjhabeng Municipality which falls under the Vaal Primary Catchment	32
Figure 3-21: Current annual and monthly surface water runoff, precipitation and evaporation for the Masilonyana Municipality which falls under the Vaal Primary Catchment	33
Figure 3-22: Projected monthly change to future (2050) evaporation, precipitation, and estimated runoff values for the Matjhabeng Municipality	34
Figure 3-23: Projected monthly change to future (2050) evaporation, precipitation, and estimated runoff values for the Masilonyana Municipality	34
Figure 3-24: Groundwater potential and dependency for the Matjhabeng and Masilonyana Municipalities (dark blue marker indicates approximate location of the project)	35
Figure 3-25: Groundwater potential and depletion for 2050 for the Matjhabeng and Masilonyana Municipalities (dark blue marker indicates approximate location of the project)	36
Figure 3-26: The contribution that the different economic sectors make to the total GVA of the Matjhabeng Municipality	37
Figure 3-27: The contribution that the different economic sectors make to the total GVA of the Masilonyana Municipality	37

LIST OF ACRONYMS AND SYMBOLS

AFF	Agriculture, Forestry and Fishing
Airshed	Airshed Planning Professionals (Pty) Ltd
AR5	IPCC's fifth assessment report
AR6	IPCC's sixth assessment report
BAU	Business-As-Usual
°C	Degrees Celsius
CAIT	Climate Analysis Indicators Tool
CCA	Climate Change Assessment
CCRA	Climate Change Reference Atlas
CCS	Carbon Capture and Sequestration (or Carbon Capture and Storage)
CH ₄	Methane
CMIP	Coupled Model Intercomparison Project
CO ₂	Carbon dioxide
CO ₂ e	Carbon dioxide equivalent
DFFE	Department of Forestry, Fisheries and the Environment (previously DEA)
DWS	Department of Water and Sanitation
EAPs	Environmental Assessment Practitioners
ECMWF	European Centre for Medium-Range Weather Forecasts
EIA	Environmental Impact Assessment
EIMS	Environmental Impact Management Services (Pty) Ltd
ER	Exploration Right
GCMs	Global Climate Change Models
GDP	Gross domestic product
GG	Government Gazette
GHG	Greenhouse gases
GHGIP	National Greenhouse Gas Improvement Programme
GN	Government Notice
GVA	Gross Value Added
GWP	Global warming potential
H ₂ O	Water vapour
HFCs	Hydrofluorocarbons
IPCC	Intergovernmental Panel on Climate Change
km	Kilometre
l/p/d	Litres per person per day
LNG	Liquefied natural gas
LUCF	Land-Use Change and Forestry
LULUCF	Land Use, Land Use Change and Forestry
m	Metre
mm	Millimetre
mm/yr	Millimetres per year
Mt	Million ton
MW	Megawatt
N ₂ O	Nitrous oxide
NAAQS	National Ambient Air Quality Standards
NAEIS	National Atmospheric Emission Inventory System
NCEP	National Centres for Environmental Prediction

NEM:AQA	National Environmental Management: Air Quality Act
NDCs	Nationally determined contributions
NOAA	National Oceanic and Atmospheric Administration
O ₃	Ozone
PFCs	Perfluorocarbons
ppm	Parts per million
PR	Production Right
PV	Photovoltaic
RCA4	Rossby Centre regional model
RCPs	Representative Concentration Pathways
SAAELIP	South African Atmospheric Emission Licensing and Inventory Portal
SAAQIS	South African Air Quality Information System
SAGERS	South African Greenhouse Gas Emission Reporting System
SAWS	South African Weather Services
SF ₆	Sulfur hexafluoride
SPI	Standardized Precipitation Index
SSP	Shared Socioeconomic Pathway
SST	Sea surface temperatures
UNFCCC	United Nations Framework Convention on Climate Change
WHO	World Health Organisation
W/m ²	Watts per metre squared
%	Percent

Note:

The spelling of "sulfur" has been standardised to the American spelling throughout the report. "The International Union of Pure and Applied Chemistry, the international professional organisation of chemists that operates under the umbrella of UNESCO, published, in 1990, a list of standard names for all chemical elements. It was decided that element 16 should be spelled "sulfur". This compromise was to ensure that in future searchable data bases would not be complicated by spelling variants. (IUPAC. Compendium of Chemical Terminology, 2nd ed. (the "Gold Book"). Compiled by A. D. McNaught and A. Wilkinson. Blackwell Scientific Publications, Oxford (1997). XML on-line corrected version: <http://goldbook.iupac.org> (2006) created by M. Nic, J. Jirat, B. Kosata; updates compiled by A. Jenkins. ISBN 0-9678550-9-8.[doi: 10.1351/goldbook](https://doi.org/10.1351/goldbook))"

Climate Change Assessment for the Motuoane Production Right Application (PR016), Free State Province – Scoping Phase

1 INTRODUCTION

Environmental Impact Management Services (Pty) Ltd (EIMS) were requested by Motuoane Energy (Pty) Ltd (Motuoane) to undertake an Environmental Impact Assessment (EIA) process for their planned new Production Right (PR) area. The proposed PR area falls within a portion of the previously assessed ER315 (exploration right). Further some of the activities being proposed by Motuoane to be included in the production right overlap with those previously assessed and authorised for the ER315 Amendment (specifically the northwestern part of the PR area). In brief the production activities will include the following:

- Provision for ongoing exploration drilling and seismic activities (preliminary seismic transects defined by the blue dashed lines in **Figure 1-1**).
- Existing wells may be ‘worked over’ and possibly included as production sites.
- Drilling and producing from approximately 30-50 production wells within the 3 identified well production corridors (as indicated in **Figure 1-1**). The intention being to assess these corridors in detail during the EIA process to allow for drilling and production within these corridors (assume a corridor width of 500 m).
- The installation of trenched sub-surface gas gathering pipelines along the red dashed lines as indicated in **Figure 1-1**.
- The pipeline network will transport gas to a local production plant located at approximately the position indicated by the red triangle in **Figure 1-1**. The gas will be compressed to deliver compressed helium and methane as a first phase followed by a second phase where the gas will be liquified to produce Liquefied Natural Gas (LNG).

Airshed Planning Professionals (Pty) Ltd (Airshed) was commissioned by EIMS to undertake a Climate Change Assessment (CCA) for the proposed Motuoane Production Right Application (PR016) (hereafter referred to as the project). This report details the scoping phase of the CCA undertaken for the project which will focus on the physical risks of climate change on the region. The climate change specialist study, which will follow the scoping phase, will assess the significance of impacts of the project greenhouse gas (GHG) emissions generated, along with the potential impact of climate change on the operation of the project.

1.1 Scope of Work

The following tasks will be included in the scope of work for the CCA scoping phase:

1. The robustness of the project in terms of forecasted climate change impacts to the area over the lifetime of the project.
2. The vulnerability of communities in the immediate vicinity of the project to climate change.
3. Potential impact of climate change in the region.




Figure 1-1: Location of the project site

2 REGULATORY CONTEXT AND IMPACT ASSESSMENT CRITERIA

GHGs are those gaseous constituents of the atmosphere, both natural and anthropogenic, that absorb and emit radiation at specific wavelengths within the spectrum of thermal infrared radiation emitted by the Earth's surface, the atmosphere itself, and by clouds. This property is known as the greenhouse effect. Water vapour (H₂O), carbon dioxide (CO₂), nitrous oxide (N₂O), methane (CH₄) and ozone (O₃) are the primary greenhouse gases in the earth's atmosphere. Moreover, there are a number of entirely human-made greenhouse gases in the atmosphere, such as the halocarbons and other chlorine and bromine containing substances, dealt with under the Montreal Protocol. Beside CO₂, N₂O and CH₄, the Kyoto Protocol deals with the greenhouse gases sulfur hexafluoride (SF₆), hydrofluorocarbons (HFCs) and perfluorocarbons (PFCs) (IPCC, 2007). Human activities since the beginning of the Industrial Revolution (taken as the year 1750) have produced a 40% increase in the atmospheric concentration of carbon dioxide, from 280 ppm in 1750 to 415 ppm in early September 2021 (NOAA, 2021). This increase of CO₂ in the Earth's atmosphere has occurred despite the uptake of a large portion of the emissions by various natural "sinks" involved in the carbon cycle (NOAA, 2021). Anthropogenic CO₂ emissions (i.e., emissions produced by human activities) come from combustion of fossil fuels, principally coal, oil, and natural gas, along with waste processing and decomposition, deforestation, soil erosion and animal agriculture (IPCC, 2007).

The following sections describe the alignment of South African national policies regarding GHG emissions and reporting with international agreements and targets.

2.1 International Agreements

In 1992, countries joined an international treaty, the United Nations Framework Convention on Climate Change, (UNFCCC) as a framework for international cooperation to combat climate change by limiting average global temperature increases and the resulting climate change, and coping with impacts that were, by then, inevitable.

By 1995, countries launched negotiations to strengthen the global response to climate change, and, two years later, adopted the Kyoto Protocol. The Kyoto Protocol legally binds developed country parties to emission reduction targets. The Protocol's first commitment period started in 2008 and ended in 2012. As agreed in Doha in 2012, the second commitment period began on 1 January 2013 and will end in 2020 (UNFCCC, 2017) but due to lack of ratification has not come into force.

The Paris Agreement (2016) builds upon the Convention and – for the first time – brings all nations into a common cause to undertake ambitious efforts to combat climate change and adapt to its effects, with enhanced support to assist developing countries to do so. As such, it charts a new course in the global climate effort.

The central aim of the Paris Agreement is to strengthen the global response to the threat of climate change by keeping a global temperature rise this century well below 2.0°C above pre-industrial levels and to pursue efforts to limit the temperature increase even further to 1.5°C. Additionally, the agreement aims to strengthen the ability of countries to deal with the impacts of climate change. To reach these ambitious goals, appropriate financial flows, a new technology framework and an enhanced capacity building framework will be put in place, thus

supporting action by developing countries and the most vulnerable countries, in line with their own national objectives.

The Paris Agreement requires all Parties to put forward their best efforts through “nationally determined contributions” (NDCs) and to strengthen these efforts in the years ahead. This includes requirements that all Parties report regularly on their emissions and on their implementation efforts.

In 2018, Parties took stock of the collective efforts in relation to progress towards the goals set in the Paris Agreement to inform the preparation of NDCs. There will also be a global stocktake every five years to assess the collective progress towards achieving the purpose of the Agreement and to inform further individual actions by Parties.

As of September 2024, 195 Parties of the 197 Parties to the UNFCCC Convention, including South Africa, had ratified the Paris Agreement. South Africa submitted its NDC to the UNFCCC on 25 September 2016 and an updated NDC in September 2021.

On January 20, 2025, an executive order was signed by the United States President to withdraw the United States from the Paris Agreement.

2.2 South African National Climate Change Response Policy

South Africa ratified the UNFCCC in August 1997 and acceded to the Kyoto protocol in 2002, with effect from 2005. However, since South Africa is an Annex 1 country it implies no binding commitment to cap or reduce GHG emissions. South Africa later also ratified the Paris Agreement (as signed on 22 April 2016) which although not bound to commit to a cap or reduce GHG emissions, pledged to reduce emissions by 34% below Business-As-Usual (BAU) emissions by 2020 and 42% below BAU by 2025. The proposed 2030 target range represents a 28% reduction in GHG emissions commitment from the original 2015 NDC targets. However, these original goals were ambitious, and South Africa subsequently shifted from BAU-based targets for 2020 and 2025 in terms of the Cancun Agreement under the UNFCCC, to absolute GHG emissions targets under the Paris Agreement. This update demonstrates reducing the upper range of South Africa’s targets by a more realistic 17% for 2025 and 28% for 2030, respectively.

The National Climate Change Response White Paper, passed by Cabinet in October 2011, stated that in responding to climate change, South Africa has two objectives: to manage the inevitable climate change impacts and to contribute to the global effort in stabilising GHG emissions at a level that avoids dangerous anthropogenic interference with the climate system. The White Paper proposes mitigation actions, especially a departure from coal-intensive electricity generation, be implemented in the short- and medium-term to match the GHG trajectory range. Peak GHG emissions are expected between 2020 and 2025 before a decade long plateau period and subsequent reductions in GHG emissions.

The White Paper also highlighted the co-benefit of reducing GHG emissions by improving air quality and reducing respiratory diseases by reducing ambient particulate matter, ozone, and sulfur dioxide concentrations to levels in compliance with the National Ambient Air Quality Standards (NAAQS) by 2020. To achieve these objectives, the

Department of Forestry, Fisheries and the Environment (DFFE) established a national GHG emissions inventory that reports through the South African Atmospheric Quality Information System (SAAQIS).

The Climate Change Act (Act 22 of 2024) was assented to by the President of the Republic of South Africa on 23 July 2024 in Government Notice (GN) 5050 in Government Gazette (GG) 50966 of 23 July 2024. The President proclaimed its commencement under section 38 on 17 March 2025; however not all provisions are in force. The Act is aligned with international policies guidelines and South Africa's NDC and aim to reduce GHG emissions as primary driver to anthropogenic climate change. The aim of the Act is to achieve an effective climate change response through a long-term just transition to a low carbon economy that is climate resilient and allows for sustainable development of South Africa. When in force, the Act will:

- establish provincial and municipal forums on climate change which will be responsible for co-ordinating climate change response actions in each province.
- strengthen the establishment of the Presidential Climate Change Coordinating Commission. Although, the commission has already been established, its establishment only carries legal force after the Bill becomes an Act.
- establish a National Adaptation Strategy to guide South Africa's adaptation to the impacts of climate change and develop adaptation scenarios which anticipate the likely impacts over the short, medium, and long term.
- determine a national GHG emissions trajectory, which must be reviewed every five years, and which indicates an emissions reduction objective.
- put in place a 5-yearly sectoral emission targets for identified sectors and sub-sectors that must be aligned with the national GHG emissions trajectory and include quantitative and qualitative GHG emission reduction goals.
- bring into force the carbon budget allocation mechanism, which will be linked to the Carbon Tax Act, which will replace the current National Pollution Prevention Plan mechanism which is enforced under the National Environmental Management: Air Quality Act (NEM:AQA).

2.3 Nationally Determined Contribution

The first South African NDC submission was completed in 2016. This was undertaken to comply with decision 1/CP.19 and 1/CP.20 of the Conference of the Parties to the UNFCCC. An update of the first NDC was published submitted to the UNFCCC on 27 September 2021¹ in preparation for the 26th Conference of the Parties (held in Glasgow, Scotland in November 2021). This document describes South Africa's NDC on adaptation, mitigation and finance and investment necessities to undertake the resolutions with updated revisions to the adaptation goals and mitigation targets.

¹

<https://www4.unfccc.int/sites/ndcstaging/PublishedDocuments/South%20Africa%20First/South%20Africa%20updated%20first%20NDC%20September%202021.pdf>

As part of the updated adaption portion the following goals have been assembled:

- Goal 1: Enhance climate change adaptation governance and legal frameworks.
- Goal 2: Develop an understanding of the impacts on South Africa of 1.5 and 2°C global warming and the underlying global emission pathways through geo-spatial mapping of the physical climate hazards, and adaptation needs in the context of strengthening the key sectors of the economy. This will provide the scientific basis for strengthening the national and provincial governments' readiness to respond to climate risk.
- Goal 3: Implementation of National Climate Change Adaptation Strategy adaptation interventions for the period 2021 to 2030, where priority sectors have been identified as biodiversity and ecosystems; water; health; energy; settlements (coastal, urban, rural); disaster risk reduction, transport infrastructure, mining, fisheries, forestry, and agriculture.
- Goal 4: Mobilise funding for adaptation implementation through multilateral funding mechanisms.
- Goal 5: Quantification and acknowledgement of the national adaptation and resilience efforts.

As part of the mitigation portion the following have been, or can be, implemented at National level:

- The approval of 79 (5 243 MW) renewable energy Independent Power Producer projects as part of a Renewable Energy Independent Power Producer Procurement Programme. An additional 6 300 MW is being deliberated.
- A "Green Climate Fund" has been created to back green economy initiatives. This fund will be increased in the future to sustain and improve successful initiatives.
- It is intended that by 2050 electricity will be decarbonised.
- Carbon Capture and Sequestration (or Carbon Capture and Storage) (CCS).
- To support the use of electric and hybrid electric vehicles.
- Reduction of emissions can be achieved through the use of energy efficient lighting; variable speed drives and efficient motors; energy efficient appliances; solar water heaters; electric and hybrid electric vehicles; solar photovoltaic (PV); wind power; CCS; and advanced bioenergy.
- Updated targets based on revised 100-year global warming potential (GWP) factors (published in the Annex to decision 18/CMA.1 of the Intergovernmental Panel on Climate Change's (IPCC) 5th assessment report) (AR5) and based on exclusion of land sector emissions arising from natural disturbance. The updated NDC mitigation targets, consistent with South Africa's fair share, are presented in **Table 2-1**.

Table 2-1: South Africa's NDC mitigation targets

Year	Target	Corresponding period
2025	South Africa's annual GHG emissions will be in a range between 398 - 510 Mt CO ₂ e.	2021-2025
2030	South Africa's annual GHG emissions will be in a range between 350 - 420 Mt CO ₂ e.	2026-2030

2.4 Greenhouse Gas Emissions Reporting

Regulations pertaining to GHG reporting using the National Atmospheric Emission Inventory System (NAEIS) were published on 3 April 2017 (Government Notice (GN) 257 in Government Gazette 40762 and amendment – GNR 994 in Government Gazette 43712). The South African mandatory reporting guidelines focus on the reporting of Scope 1 emissions only. The three broad scopes for estimating GHG are:

- Scope 1: All direct GHG emissions.
- Scope 2: Indirect GHG emissions from consumption of purchased electricity, heat or steam.
- Scope 3: Other indirect emissions, such as the extraction and production of purchased materials and fuels, transport-related activities in vehicles not owned or controlled by the reporting entity, electricity-related activities not covered in Scope 2, outsourced activities, waste disposal, etc.

The South African Greenhouse Gas Emission Reporting System (SAGERS) web-based monitoring and reporting system is used to collect GHG information in a standard format for comparison and analyses. The system forms part of the national atmospheric emission inventory component of South African Atmospheric Emission Licensing and Inventory Portal (SAAELIP).

The DFFE is working together with local sectors to develop country specific emissions factors in certain areas; however, in the interim the IPCC's default emission figures may be used to populate the SAAQIS GHG emission factor database. These country specific emission factors will replace some of the default IPCC emission factors. Methodological guidelines for GHG emission estimation, which include country specific emission factors for fuels used in stationary and mobile combustion, have been issued (DFFE, 2022b).

Also, the Carbon Tax Act (Act 15 of 2019) includes details on the imposition of a tax on the carbon dioxide equivalent (CO₂e) of GHG emissions. Certain production processes indicated in Annexure A of the Declaration of Greenhouse Gases as Priority Pollutants (GN 710 in GG 40966, 21 July 2017) with GHG more than 0.1 Mt/year, measured as CO₂e, are required to submit a greenhouse gas mitigation plan to the Minister for approval. The proposed project will be required to report CO₂e emissions and to prepare a greenhouse gas mitigation plan.

2.5 GHG Inventories

2.5.1 National GHG Emissions Inventory

South Africa is a GHG contributor and is undertaking steps to mitigate and adapt to the changing climate. DFFE is categorised as the lead climate change institution and is required to coordinate and manage climate related information such as development of mitigation, monitoring, adaption and evaluation strategies (DFFE, 2022a). This includes the establishment and updating of the National GHG Inventory. The National Greenhouse Gas Improvement Programme (GHGIP) has been initiated; it includes sector specific targets to improve methodology and emission factors used for the different sectors as well as the availability of data.

The 2020 National GHG Inventory was prepared using the 2006 IPCC Guidelines (IPCC, 2006). According to the 9th National GHG Inventory Report (DFFE, 2025), the total GHG emissions in 2022 were estimated at

approximately 478.3009 Mt CO₂e (excluding Land Use, Land Use Change and Forestry (LULUCF)). This was a 2.2% decrease from the 2000 total GHG emissions (excluding LULUCF). LULUCF is estimated to be a net carbon sink which reduces the 2022 GHG emissions to 435.1196 Mt CO₂e. The assessment (excluding LULUCF) showed the main sector contributing to GHG emissions in 2022 to be the energy sector, contributing 78% to the total GHG emissions.

2.5.2 GHG Emission Inventory for the Sector

The proposed project would be categorised in the energy category for both the global GHG inventory and for the national GHG inventory. According to the World Resources Institute – Climate Analysis Indicators Tool (CAIT) Climate Data Explorer² the 2022 global GHG emissions from the energy category were approximately 38 008 Mt CO₂e; 75.8% of the total GHG emissions (including Land-Use Change and Forestry (LUCF)). The South African energy sector contributed 374.1141 Mt CO₂e, ~1% of the global emissions from the energy sector in 2022.

2.5.3 Draft National Guideline for Consideration of Climate Change in Development Applications, June 2021

The DFFE published (on 25 June 2021) a notice under the NEMA requesting public comment on the *Draft National Guideline for the consideration of climate change implications in applications for environmental authorisation, atmospheric emission licences and waste management licences*.

The Draft National Guideline has been developed to support the inclusion of climate change considerations into the Environmental Impact Assessment (EIA) process, and to create a consistent approach for such incorporation, which will help proponents to assess:

- how a proposed development will likely exacerbate climate change;
- the impact of a development on features (natural and built) that are crucial for climate change adaptation and resilience; and,
- the sustainability of a development in the context of climate change projection.

The Guideline puts forward “a consistent approach in providing interested and affected parties (for example, proponents, Environmental Assessment Practitioners (EAPs) and specialists) with the minimum requirements to consider when undertaking a climate change assessment, which forms part of an application for environmental authorisation, an atmospheric emissions licence, and/or waste management licence”.

One of the impact requirements for a climate change assessment is an estimation of the GHG emissions, direct and indirect (including upstream GHG emissions) that will be released into the atmosphere annually throughout the impact related to the activity.

² <http://cait.wri.org/>

The comment period for amendments to the draft guideline has now closed but the final guideline has not yet been published. As far as possible the guideline has been followed in the preparation of this climate change impact assessment in support of environmental authorisation.

3 PHYSICAL RISKS OF CLIMATE CHANGE ON THE REGION

The discussions of physical risks of climate change discussed in this section are likely to be relevant to the project as well as to the communities surrounding the project even if the project is not authorised.

3.1 Vulnerability

The Green Book (CSIR, 2025); was developed to be an online platform providing quantitative scientific evidence on the likely impacts that climate change and urbanisation will have on South Africa's cities and towns. A profile for each local municipality, including individual settlements and neighbourhoods, was built in terms the rates of socio-economic, economic, physical and environmental risks associated with urbanisation, population growth and climate change (Le Roux, et al., 2019). The risk profile was accessed for the Matjhabeng and Masilonyana Municipalities³.

The Matjhabeng Municipality socio-economic vulnerability score⁴ (out of 10) is 5.3 for 1996, reducing to 4.2 for 2011. The lower score in 2011 compared to 1996 indicates improvement of socio-economic factors. The Matjhabeng Municipality for socio-economic vulnerability ranks 4th out of 19 in the province and 81st out of 213 in the country. The Matjhabeng Municipality economic vulnerability score⁵ (out of 10) is 7.7 for 1996, increasing to 9.9 for 2011. The economic vulnerability ranks 19th out of 19 in the province and 211th out of 213 in the country. The physical vulnerabilities⁶ ranks 4th out of 19 in the province and 50th out of 213 in the country. The environmental vulnerability⁷ ranks 13th out of 19 in the province and 102nd out of 213 in the country.

The Masilonyana Municipality socio-economic vulnerability score (out of 10) is 6.7 for 1996, reducing to 5.8 for 2011. The lower score in 2011 compared to 1996 indicates improvement of socio-economic factors. The Masilonyana Municipality for socio-economic vulnerability ranks 15th out of 19 in the province and 141st out of 213 in the country. The Masilonyana Municipality economic vulnerability score (out of 10) is 7.4 for 1996, increasing to 8.7 for 2011. The economic vulnerability ranks 18th out of 19 in the province and 202nd out of 213 in the country. The physical vulnerabilities ranks 6th out of 19 in the province and 57th out of 213 in the country. The environmental vulnerability ranks 2nd out of 19 in the province and 31st out of 213 in the country.

³ <https://riskprofiles.greenbook.co.za/>

⁴ Defined as the vulnerability of households based on household composition; education and health; access to basic services; safety and security.

⁵ Defined as the susceptibility of the municipality to external shocks based on economic diversity; size of economy; labour force; gross domestic product (GDP) growth rate; and inequality.

⁶ Defined by the physical fabric of connectedness of the settlements within the municipalities and structural robustness.

⁷ This indicator represents the balance between preserving the natural environmental and the pressures of population growth, urbanisation, and economic development. The indicator is based on air quality, environmental governance and competition between ecology and the urban environment.

3.2 Climate

3.2.1 Baseline Climate

Climate change metrics focus on temperature; the number of very hot days (where temperatures exceed 35°C); and rainfall. The baseline (1961 to 1990) annual averages for these metrics were accessed for the area near the project site from the South Africa 'Green Book'⁸ (CSIR, 2025).

Baseline annual average temperature was in the range 15.9°C (**Figure 3-1**) with the three very hot days per year and seven heat wave days per year (**Figure 3-2**). Average rainfall (**Figure 3-1**) was 1 099 mm.

⁸ <https://greenbook.co.za/>




Figure 3-1: Baseline (1961 to 1990) annual average temperature and rainfall for the project area (CSIR, 2025) (dark blue marker indicates approximate location of the project)




Figure 3-2: Baseline (1961 to 1990) number of very hot days (>35°C) and heat wave days¹ annually for the project area (CSIR, 2025) (dark blue marker indicates approximate location of the project)

¹ The annual heat-wave days map depicts the number of days (per 8x8 km grid point) where the maximum temperature exceeds the average maximum temperature of the warmest month of the year at that location by 5°C, for a period of at least three consecutive days.

Recent change in climatic conditions near the project site were accessed from MeteoBlue¹ a weather forecasting platform developed at the University of Basel, Switzerland and based on models of National Oceanic and Atmospheric Administration (NOAA) or National Centres for Environmental Prediction (NCEP). The data sets also include historical climate data tracking changes in climate by referencing ERA5, the fifth generation ECMWF (European Centre for Medium-Range Weather Forecasts) atmospheric reanalysis of the global climate, for the period between 1979 to 2024, with a spatial resolution of 30 km. Based on the study area, an increasing trend in the annual average temperatures has been observed with temperatures measuring 16.7°C in 1979 to 18.1°C in 2024 (**Figure 3-3** – top panel). The lower part the graph shows the so-called warming stripes. Each coloured stripe represents the average temperature for a year - blue for colder and red for warmer years. The change in rainfall over the same period (1979 – 2024) displays a slight decreasing trend (Figure 3-6), where the difference from long-term average for each year in the data set is visualised by the stripes in the lower panel of **Figure 3-4** (brown stripes indicate lower than average rainfall and green stripes above average rainfall).




Figure 3-3: Annual average temperature (top panel) and temperature anomaly (lower panel) between 1979 and 2024 (Meteoblue, 2025)

¹ <https://www.meteoblue.com>




Figure 3-4: Annual average rainfall (top panel) and rainfall anomaly (lower panel) between 1979 and 2024 (Meteoblue, 2025)

3.2.2 Projected Future Climate

In 2017 the South African Weather Services (SAWS) published an updated Climate Change Reference Atlas (CCRA) based on Global Climate Change Models (GCMs) projections (SAWS, 2017). It must be noted that as with all atmospheric models there is the possibility of inaccuracies in the results because of the model's physics and accuracy of input data. The Rossby Centre regional model (RCA4) was used in the predictions for the CCRA which included the input of nine GCMs results. The RCA4 model was used to improve the spatial resolution to $0.44^{\circ} \times 0.44^{\circ}$ - the finest resolution GCMs in the ensemble were run at resolutions of $1.4^{\circ} \times 1.4^{\circ}$ and $1.8^{\circ} \times 1.2^{\circ}$. Findings from downscaled climatic simulations using six GCMs, at an 8 km x 8 km resolution over South Africa, for the time slab 2021 to 2050 were included in the Green Book (Engelbrecht, 2019).

In both the CCRA and the Green Book, two trajectories are included based on the four Representative Concentration Pathways (RCPs) discussed in the IPCC's fifth assessment report (AR5) (IPCC, 2013). RCPs are defined by their influence on atmospheric radiative forcing in the year 2100. RCP4.5 represents an addition to the radiation budget of 4.5 W/m^2 as a result of an increase in GHGs. The two RCPs selected were RCP4.5 representing the medium-to-low pathway and RCP8.5 representing the high pathway. RCP4.5 is based on a CO_2 concentration of 560 ppm and RCP8.5 on 950 ppm by 2100. RCP4.5 is based on if current interventions to reduce GHG emissions being sustained (after 2100 the concentration is expected to stabilise or even decrease). RCP8.5 is based on if no interventions to reduce GHG emissions being implemented (after 2100 the concentration is expected to continue to increase).

3.2.2.1 RCP4.5 Trajectory

The Green Book projected temperature changes in the near future (up to 2050) indicate an average increase of 2.3°C (**Figure 3-5**, Engelbrecht, et al., 2019). The number of very hot days are expected to increase by 8.5 days per year (**Figure 3-6**). Between 2021 and 2050 the annual rainfall near the project site was projected to increase by 163 mm per year (**Figure 3-7**, Engelbrecht, et al., 2019), with extreme rainfall days potentially increasing by 2.2 days in the near future (**Figure 3-8**, Engelbrecht, et al., 2019).




Figure 3-5: Projected change in annual average temperature for the near future (2021 – 2050) (dark blue marker indicates approximate location of the project)




Figure 3-6: Projected change in annual average number of very hot days for the near future (2021 – 2050) (dark blue marker indicates approximate location of the project)




Figure 3-7: Projected change in annual average rainfall for the near future (2021 – 2050) (dark blue marker indicates approximate location of the project)




Figure 3-8: Projected change in annual average number of extreme rainfall days (>20 mm in <24 hours) for RCP4.5 trajectory (dark blue marker indicates approximate location of the project)

3.2.2.2 RCP8.5 Trajectory

The Green Book projected temperature changes in the near future (up to 2050) indicate an average increase of 2.6°C (**Figure 3-5**, Engelbrecht, et al., 2019). The number of very hot days are expected to increase to between 9 and 34 days per year. Between 2021 and 2050 the annual rainfall near the project site was projected to increase by 226 mm per year (**Figure 3-7**, Engelbrecht, et al., 2019), with extreme rainfall days potentially increasing by 2.1 days in the near future (**Figure 3-8**, Engelbrecht, et al., 2019).

3.2.2.3 IPCC's Sixth Assessment Report: Temperature and Rainfall Projections

The most recent IPCC data are from the Coupled Model Intercomparison Project (CMIP) which were derived from the sixth phase of the CMIPs (CMIP6) and supports the IPCC's Sixth Assessment Report (AR6) which was released on 9 August 2021 (Working Group I), 28 February 2022 (Working Group II and 4 April 2022 (Working Group III). Projection data is presented at a 1.0° x 1.0° (100 km x 100 km) resolution. The scenarios are the result of complex calculations that depend on how quickly humans curb greenhouse gas emissions, whilst also capturing socioeconomic changes in areas such as population, urban density, education, land use and wealth. For example, a rise in population is assumed to lead to higher demand for fossil fuels and water. Education can affect the rate of technology developments. Emissions increase when land is converted from forest to agricultural land. Each scenario is labelled to identify both the emissions level and the so-called Shared Socioeconomic Pathway, or SSP, used in those calculations. This first scenario is the only one that meets the Paris Agreement's goal of keeping global warming to around 1.5°C above preindustrial temperatures, with warming hitting 1.5°C but then dipping back down and stabilizing around 1.4°C by the end of the century. Projected changes are defined relative to a historical 20-year period (1995 to 2014).

The AR6 projections for the study area for the scenario RCP4.5 indicate an increase in annual average temperatures of 1.6°C for the period 2041 to 2060 and 2.3°C for the period 2081 to 2100. The projections for the RCP8.5 indicate an increase in annual average temperatures of 2.7°C for the period 2041 to 2060, to 5.3°C for the period 2081 to 2100 (IPPC, 2022). The AR5 projections, for comparison, estimate an increase in annual average temperatures (50th percentile) of 2.3°C for RCP4.5 and 2.6°C for RCP8.5 for the period 2021 to 2050.

The AR6 projections for rainfall in the study area for RCP4.5 indicate a decrease in annual rainfall of 5% for the period 2041 to 2060, to 6% for the period 2081 to 2100. The projections for RCP8.5 indicate a decrease in rainfall of 6% for the period 2041 to 2060, to 10% for the period 2081 to 2100 (IPPC, 2022).

3.3 Hazards

The Green Book risk profile includes an assessment of projected risk to the Matjhabeng and Masilonyana Municipalities in 2050, mostly based on the low mitigation RCP8.5 climate simulations, and highlights the following:

- Isolated pockets of moderate and high increased risk of wildfires within the municipalities, but not within the project area (**Figure 3-9**);
- Isolated pockets of very low risk of increased drought frequency within the municipalities and the Standardized Precipitation Index (SPI)¹ of -0.44 (**Figure 3-10**);
- Isolated pockets of high increased heat extremes within the municipalities (**Figure 3-11**); and,
- Areas of slight to moderate risk of increased extreme rainfall days within the project area (**Figure 3-12**).
- Isolated pockets of very low to moderate increase in exposure to urban flooding is expected for the municipalities (**Figure 3-12**).




Figure 3-9: Risk of increased wildfires for the Matjhabeng and Masilonyana Municipalities in 2050 based on RCP8.5 trajectory (dark blue marker indicates approximate location of the project)

¹ The Standardized Precipitation Index (SPI) is a widely used index to characterize meteorological drought on a range of timescales. SPI index.




Figure 3-10: Risk of increased drought tendencies for the Matjhabeng and Masilonyana Municipalities in 2050 based on RCP8.5 trajectory (dark blue marker indicates approximate location of the project)




Figure 3-11: Risk of increased heat extremes for the Matjhabeng and Masilonyana Municipalities in 2050 based on RCP8.5 trajectory (dark blue marker indicates approximate location of the project)




Figure 3-12: Risk of increased flooding for the Matjhabeng and Masilonyana Municipalities in 2050 based on RCP8.5 trajectory (dark blue marker indicates approximate location of the project)

In addition to the hazards identified in the Green Book, Hofste, *et al.*, (2019) currently rate the project area as arid and low water use (Figure 3-13) with a near normal water stress projection for the future (2050 based on a conservative low mitigation trajectory) (Figure 3-14).




Figure 3-13: Current water stress for the project area (Hofste, et al., 2019) (blue dot indicates project location)




Figure 3-14: Projected (2050) water stress for the project area (Hofste, et al., 2019) (blue dot indicates project location)

3.4 Vegetation Disturbance

The potential vegetation types that will be disturbed over the lifetime of the proposed project include (South African National Biodiversity Institute, 2025):

- Bloemfontein Karroid Shrubland;
- Central Free State Grassland;
- Eastern Free State Clay Grassland;
- Highveld Alluvial Vegetation; and,
- Vaal-Vet Sandy Grassland.

This information will be used in the quantification of permanent GHG sink losses due to project infrastructure for the CCA.

3.5 Impact of Climate Change

To understand the impact that climate change might have on the major resources of the Matjhabeng and Masilonyana Municipalities it is first necessary to provide an overview of the current situation, which has been provided for water, economy, and agriculture.

3.5.1 Water Supply

3.5.1.1 Current Resources

Figure 3-15 provides the current water supply vulnerability (i.e., demand versus supply) for the Matjhabeng Municipality (1.43) based on the data compiled for the Department of Water and Sanitation (DWS) All Town's Study (Cole, Bailey, Cullis, & New, 2017). The current water demand for the municipality is 207 l/p/d (litres per person per day) with supply of 144.5 l/p/d, with 100% sourced from surface water.

Figure 3-16 provides the current water supply vulnerability (i.e., demand versus supply) for the Masilonyana Municipality (1) based on the data compiled for the Department of Water and Sanitation (DWS) All Town's Study (Cole, Bailey, Cullis, & New, 2017). The current water demand for the municipality is 257 l/p/d (litres per person per day) with supply of 258 l/p/d, with 80% sourced from surface water and 20% from groundwater.




Figure 3-15: Current water availability for the Matjhabeng Municipality





Figure 3-16: Current water availability for the Masilonyana Municipality

3.5.1.2 Impact on Resources

Figure 3-17 and **Figure 3-18** shows the estimated current and future water supply vulnerability (i.e., the ratio of demand to supply) based on: 1) a local water supply perspective incorporating changes to population growth coupled with exposure to climate risk (based on impacts on local runoff), and 2) a regional water supply perspective (based on impacts of regional water supply assuming supply is part of the integrated regional and national bulk water supply network). The mean annual precipitation for the Matjhabeng and Masilonyana municipalities is predicted to increase by 6.5% and 6.9% respectively for 2050 with a regional urban water supply increase of 11.9%.














VULNERABILITY CONTRIBUTION FACTORS			PERCENTAGE CHANGE	
	Mean annual precipitation		6.5%	
	Mean annual evaporation		8.02%	
	Mean annual runoff		56.82%	
	Regional urban water supply		11.91%	
	Population growth		-4.72%	

Figure 3-17: Estimated current and future (2050) water supply vulnerability based on medium population growth for the Matjhabeng Municipal







VULNERABILITY CONTRIBUTION FACTORS			PERCENTAGE CHANGE
	Mean annual precipitation		6.92%
	Mean annual evaporation		8.03%
	Mean annual runoff		42.72%
	Regional urban water supply		11.91%
	Population growth		-1.95%

Figure 3-18: Estimated current and future (2050) water supply vulnerability based on medium population growth for the Masilonyana Municipal

3.5.2 Surface Water

3.5.2.1 Current Situation

The project location is within the Vaal Primary Catchment (**Figure 3-19**).

Figure 3-20 and **Figure 3-21** depicts the current annual and monthly surface water runoff, precipitation and evaporation for the Vaal Primary Catchment associated with the Matjhabeng and Masilonyana municipalities respectively. Precipitation and evaporation for the Matjhabeng Municipality is currently 533 mm/yr and 1 615 mm/yr respectively. Precipitation and evaporation for the Masilonyana Municipality is currently 525 mm/yr and 1 612 mm/yr respectively.




Figure 3-19: Quaternary catchment areas for the study area




Figure 3-20: Current annual and monthly surface water runoff, precipitation and evaporation for the Matjhabeng Municipality which falls under the Vaal Primary Catchment




Figure 3-21: Current annual and monthly surface water runoff, precipitation and evaporation for the Masilonyana Municipality which falls under the Vaal Primary Catchment

3.5.2.2 Projected Impact

Figure 3-22 and **Figure 3-23** provides the projected monthly change for future (2050) evaporation, precipitation, and estimated runoff values for the Matjhabeng and Masilonyana municipalities respectively.




Figure 3-22: Projected monthly change to future (2050) evaporation, precipitation, and estimated runoff values for the Matjhabeng Municipality




Figure 3-23: Projected monthly change to future (2050) evaporation, precipitation, and estimated runoff values for the Masilonyana Municipality

3.5.3 Ground Water

3.5.3.1 Current Situation

The groundwater recharge potential map indicates the occurrence and distribution of groundwater resources across the country, showing distinctive recharge potential zones. The groundwater dependency map indicates where settlements get their main water supply from, be it groundwater, surface water or a combination of both sources. Settlements that rely on groundwater, either entirely or partially, are deemed groundwater dependent. The residential settlements within the vicinity of the project area are surface water dependent.




Figure 3-24: Groundwater potential and dependency for the Matjhabeng and Masilonyana Municipalities (dark blue marker indicates approximate location of the project)

3.5.3.2 Projected Impact

A groundwater depletion risk map was created to determine which of South Africa's groundwater dependent settlements may be most at risk to groundwater depletion based on decreasing groundwater aquifer recharge potential and significant increases in population growth pressure by 2050. The groundwater depletion risk map (**Figure 3-25**) is based on the settlement aquifer recharge potential of the 50th percentile RCP8.5 scenario, and the medium population growth scenario. Based on this information, and based on the fact that the nearest

residential areas are not groundwater dependent, there is no groundwater depletion risk within the areas closest to the study area.




Figure 3-25: Groundwater potential and depletion for 2050 for the Matjhabeng and Masilonyana Municipalities (dark blue marker indicates approximate location of the project)

3.5.4 Economy

Figure 3-26 and **Figure 3-27** shows the contribution that the different economic sectors make to the total Gross Value Added (GVA)¹² of the Matjhabeng Municipality and the Masilonyana Municipality respectively as well as its national GVA rank (total GVA contribution to the national GVA). The Matjhabeng Municipality ranks 15th in the national GVA rank with mining and quarrying activities making up the highest economic sector to the total GVA at 41.1%. The Masilonyana Municipality ranks 128th in the national GVA rank with mining and quarrying activities making up the highest economic sector to the total GVA at 38.6%.

¹² Gross value added (GVA) is an economic productivity metric that measures the contribution of a corporate subsidiary, company, or municipality to an economy, producer, sector, or region.




Figure 3-26: The contribution that the different economic sectors make to the total GVA of the Matjhabeng Municipality




Figure 3-27: The contribution that the different economic sectors make to the total GVA of the Masilonyana Municipality

Table 3-1 and **Table 3-2** summarises the forecasted economic gains or losses for the Matjhabeng Municipality and Masilonyana Municipality respectively, under both the RCP4.5 and RCP8.5 scenarios, for each of the contributing economic sectors.

Table 3-1: Forecasted economic gains or losses for the Matjhabeng Municipality for the RCP4.5 and RCP8.5 scenarios

RCP 4.5 Impacts			RCP 8.5 Impacts		
Average	0.92%	▲	Average	-0.05%	▼
Agriculture Sector	1.23%	▲	Agriculture Sector	-0.64%	▼
Forestry Sector	1.73%	▲	Forestry Sector	-0.91%	▼
Fishing Sector	1.24%	▲	Fishing Sector	-0.65%	▼
Mining Sector	-0.1%	▼	Mining Sector	-0.03%	▼
Manufacturing Sector	-0.54%	▼	Manufacturing Sector	0.76%	▲
Electricity & Gas Sector	4.78%	▲	Electricity & Gas Sector	1.29%	▲
Water Sector	-1.54%	▼	Water Sector	-0.42%	▼
Service Sector	0.57%	▲	Service Sector	0.17%	▲

Table 3-2: Forecasted economic gains or losses for the Masilonyana Municipality for the RCP4.5 and RCP8.5 scenarios

RCP 4.5 Impacts			RCP 8.5 Impacts		
Average	-3.97%	▼	Average	-4.6%	▼
Agriculture Sector	-7.21%	▼	Agriculture Sector	-8.45%	▼
Forestry Sector	-16.56%	▼	Forestry Sector	-19.4%	▼
Fishing Sector	0%	-	Fishing Sector	0%	-
Mining Sector	0.28%	▲	Mining Sector	0.33%	▲
Manufacturing Sector	2.91%	▲	Manufacturing Sector	3.85%	▲
Electricity & Gas Sector	-13.61%	▼	Electricity & Gas Sector	-15.95%	▼
Water Sector	4.07%	▲	Water Sector	4.77%	▲
Service Sector	-1.67%	▼	Service Sector	-1.95%	▼

3.5.5 Agriculture, Forestry and Fisheries

The main agricultural commodities for the Matjhabeng Municipality are maize, beef cattle and milk and cream (**Table 3-3**). Agriculture, Forestry and Fishing (AFF) sector contributes 1.12% to Matjhabeng Municipality GVA production and 4.28% to Matjhabeng total employment. The total AFF GVA production of Matjhabeng Municipality contributes 0.5% to the national AFF GVA (**Table 3-3**).

The main agricultural commodities for the Masilonyana Municipality are maize and beef cattle (**Table 3-4**). Agriculture, Forestry and Fishing (AFF) sector contributes 6.8% to Masilonyana Municipality GVA production and 21.1% to Masilonyana total employment. The total AFF GVA production of Masilonyana Municipality contributes 0.25% of the national AFF GVA (**Table 3-4**).

Table 3-3: Economic contribution of main commodities for Matjhabeng Municipality




MAIN COMMODITIES		
 MAIZE FOR GRAIN	 BEEF CATTLE	 MILK AND CREAM
AFF contributes 1.12% to Matjhabeng GVA production	AFF contributes 4.28% to Matjhabeng total employment	The total AFF GVA production of Matjhabeng Municipality contributes 0.51% to the national AFF GVA, ranking them as the 62nd biggest contributor

Table 3-4: Economic contribution of main commodities for Masilonyana Municipality

MAIN COMMODITIES		
 BEEF CATTLE	 MAIZE FOR GRAIN	
AFF contributes 6.76% to Masilonyana GVA production	AFF contributes 21.14% to Masilonyana total employment	The total AFF GVA production of Masilonyana Municipality contributes 0.25% to the national AFF GVA, ranking them as the 118th biggest contributor

For the Matjhabeng Municipality, the main agricultural commodities for 2050 are maize, beef cattle and milk and cream (under an RCP8.5 low-mitigation scenario) (**Table 3-5**). The climate for the municipality is expected to be hotter and wetter with slightly more extreme rainfall events. There is the potential increase in maize yield for the near future. However, towards 2050, heat stress can negatively impact on production. The hot moist conditions will result in increased spread of disease and parasites, and the heat stress will result in reduced growth and reproduction performance for beef cattle and milk yield and milk quality.

For the Masilonyana Municipality, the main agricultural commodities for 2050 are maize and beef cattle (under an RCP8.5 low-mitigation scenario) (**Table 3-6**). The climate for the municipality is expected to be hotter and wetter with more extreme rainfall events. There is the potential increase in maize yield for the near future. However, crop suitability may decline over time as temperatures start to exceed critical crop thresholds. The hot moist conditions will result in increased spread of disease and parasites, and the heat stress will result in reduced growth and reproduction performance for beef cattle.

Table 3-5: Projected economic contribution of main commodities for Matjhabeng Municipality






MAIN COMMODITIES		
 <p>MAIZE FOR GRAIN</p>	 <p>BEEF CATTLE</p>	 <p>MILK AND CREAM</p>
CLIMATE IMPACT		
<p>Change in climate expected: Hotter and wetter with more extreme rainfall events.</p>		
<p>Potential increase in maize yield for near future. However, towards 2050, heat stress can negatively impact on production.</p>	<p>Increased water availability. Hot and moist conditions cause increased spread of disease and parasites. Reduced growth & reproduction performance due to heat stress.</p>	<p>Hot and moist conditions cause increased spread of disease and parasites. Potential increase in heat stress which could negatively affect conception rates, milk yield and milk quality.</p>

Table 3-6: Projected economic contribution of main commodities for Masilonyana Municipality

MAIN COMMODITIES	
 <p>BEEF CATTLE</p>	 <p>MAIZE FOR GRAIN</p>
CLIMATE IMPACT	
<p>Change in climate expected: Hotter and wetter with more extreme rainfall events.</p>	
<p>Increased water availability. Hot and moist conditions cause increased spread of disease and parasites. Reduced growth & reproduction performance due to heat stress.</p>	<p>Potential increase in maize yield for near future. However, towards 2050, heat stress can negatively impact on production.</p>

3.5.6 Other Resources

The impacts of climate change on other resources are summarised in **Table 3-7**.

Table 3-7: The impacts of climate change on other resources

Parameter	Results of Climate Change				Reference
	Increase in temperature and heat stress	Drought and decrease in rainfall	Increase in rainfall and inland flooding	Increased wind speed	
Transport and Mobility	<ul style="list-style-type: none"> Increased rate of infrastructure deterioration leading to pavement failure including cracking, rutting, potholes, flushing, and stripping. Increased stress on bridges, particularly expansion joints, through thermal expansion and increased movement. Corrosion of steel reinforcing in concrete structures due to increase in surface salt levels in some locations. Increased infrastructure maintenance cost for road repair and reconstruction work, causing traffic delays and emergency service response delays. Increased frequency and intensity of wildfires leading to more road closures. Increased vehicle accidents, due to low pavement adhesion, leading to higher rates of transport-related fatalities. 	<ul style="list-style-type: none"> Reduced water resources available for construction and maintenance. Reduced production of some agricultural produce leading to changes in freight flows in the network. 	<ul style="list-style-type: none"> Increased rate of infrastructure deterioration, especially in areas with poor infrastructure maintenance history. Temporary and permanent flooding of road, rail, port and airport infrastructure. Structural integrity of roads, bridges and tunnels could be compromised by higher soil moisture levels. Potential destruction of bridges and culverts. Erosion of embankments and road bases leading to undermining of roads or railways. Increased risk of landslides, slope failures, road washouts and closures. Undermining of bridge structures (scouring). Closure of roadways and tunnels leading to traffic delays. Transportation system disruptions, impacts to traffic signalling and low water crossings. Increased weather-related accidents. 	<ul style="list-style-type: none"> Increased drag on vehicles resulting in increased fuel consumption. Increased safety risk for pedestrians and cyclists due to flying objects or being uncontrollably dragged by winds, additionally leading to reduced trip making by pedestrians and cyclists. 	(Mokonyama & Van Wyk, 2018)
Solid Waste	<ul style="list-style-type: none"> Increased risk of combustion at open waste disposal sites and illegal dumps and increase in explosion risk associated with methane gas. Increased rate of decay of putrescible waste resulting in increased odour, breeding of flies, and attracting of vermin. Increased health and safety concern regarding heat stroke to staff collecting waste. Increased risk of landfill site instability and failure due to changes in consumption patterns with increased waste creation (i.e., glass, plastic and paper cups). 		<ul style="list-style-type: none"> Increased risk of flooding due to pressure on stormwater and leachate management systems at landfills. Increased demand for capacity to cope with large volumes of waste generated by flood events. Increase in soil saturation causing decreased stability of slopes and landfills linings (if clay or soil based) at waste management facilities. Inundation of waste releasing contaminants to waterways, pathways and low elevation zones. Potential loss of value and degradation of paper and cardboard for recycling due to increased moisture content. Increased flooding causing the risk of localised disruption of waste collection rounds. Flooding in areas with untreated, dumped waste causing the risk of groundwater contamination. Increased flooding causing the risk of litter entering the storm water systems. 	<ul style="list-style-type: none"> Possible increase in nuisance due to waste dispersed by high winds leading to increased health effects associated with particulate matter (air pollution). 	(Oelofse, 2018)

Parameter	Results of Climate Change				Reference
	Increase in temperature and heat stress	Drought and decrease in rainfall	Increase in rainfall and inland flooding	Increased wind speed	
Stormwater	<ul style="list-style-type: none"> • Potential risk of undermining the temperature regime of temperature-sensitive stormwater ponds and receiving waters, resulting in a decrease in water quality. • Increased corrosion in stormwater drains due to a combination of higher temperatures, increased strengths, longer retention times, and stranding of solids. 	<ul style="list-style-type: none"> • Increased shrinking soils increasing the potential for cracking, increased infiltration and exfiltration of water mains and sewers, which in turn exacerbates treatment and groundwater or storm water contamination. 	<ul style="list-style-type: none"> • Increased risk of flooding due to pressure on stormwater systems. • Increased risk of litter entering the stormwater systems. • Increased risk of damage and failure of stormwater systems due to overloading during floods and intense rainfall events. • Failure of stormwater treatment devices during high flow events leading to by-pass and / or flushing of contaminated water. • High wet-weather hydraulic loads and bottlenecks in stormwater and networks due to inflow and sewer infiltration, leading to local inundation and overflows of untreated wastewater. • Increased rainfall causes soil erosion thus damaging underground stormwater systems. • Increased surface and stream erosion causing deposition of sediments in receiving environments. • Stream morphology for undeveloped, developing and fully developed urban areas, may change, hence affecting existing outfall structures and potential stormwater pond locations. 	<ul style="list-style-type: none"> • Increased wind speed and intensity causing changes in rainfall over complex topography including increasing upwind of hills and ranges. 	(Dunker & Van Wyk, 2018)
Sanitation	<ul style="list-style-type: none"> • Increased heat waves, accompanied by dry weather, can exacerbate already stressed water supply systems leading to competition between sectors for water services, affecting sanitation. 	<ul style="list-style-type: none"> • Decrease in water supply for sanitation through decrease in available water to flush sewage systems adequately. • Declining annual rainfall threatening the viability of water-borne sanitation systems, and the capacity of surface water to dilute, attenuate and remove pollution. • Sewers are structurally vulnerable to drying, hence shrinking soils increase the potential for cracking, increased infiltration, and exfiltration, which in turn exacerbates treatment and groundwater or storm water contamination. • Increased corrosion in sewers due to a combination of higher temperatures, increased strengths, longer retention times, and stranding of solids. 	<ul style="list-style-type: none"> • Increased wet-weather hydraulic loads and bottleneck in stormwater and sanitary sewer networks due to inflow and sewer infiltration, causing local inundation and overflows of untreated wastewater. • Increased rainfall and heavy rainfall events increasing the washing of faecal matter into water sources due to flooding of wastewater treatment works. • Increased risk of flooding resulting in both infrastructure damage and contamination of surface and groundwater supplies. • Increased groundwater levels due to flooding, putting risk on sewage treatment plants (which are often positioned on low-lying ground as sewerage systems rely on gravity). • Increased vulnerability of sewerage pipe systems due to their size and complexity, and their exposure to multiple flood damage threats from source, through treatment, to delivery. • Increased vulnerability of pit toilets (widely used in rural areas) due to flooding, causing serious environmental contamination. • Increase in groundwater recharge and groundwater levels causing flooding of subsurface infrastructure such as pit toilets or septic tanks. 		(Duncker, 2018)

Parameter	Results of Climate Change				Reference
	Increase in temperature and heat stress	Drought and decrease in rainfall	Increase in rainfall and inland flooding	Increased wind speed	
Information and Communication Technology	<ul style="list-style-type: none"> Increased weathering and deterioration of infrastructure resulting in increased maintenance and repair costs. Heat stress causing structural damage to infrastructure. Increased energy demands during heatwaves resulting in power outages which can impact on delivery of telecommunications services. Increases in temperature and higher frequency, duration, and intensity of heat waves increasing the risk of overheating in data centres, exchanges, and base stations, which can result in increased failure rates of equipment. Increased mean temperature increasing operating temperature of network equipment which may cause malfunctions if it surpasses design limits. 	<ul style="list-style-type: none"> Decreased precipitation leading to land subsidence and heave, reducing the stability of telecommunications infrastructure above and below ground (foundations and tower structures). 	<ul style="list-style-type: none"> Increased risk of flooding of low-lying infrastructure, access holes and underground facilities. Increases in storm frequency or intensity increasing the risk of damage to aboveground transmission infrastructure and impacting on telecommunications service delivery. Increases in storm frequency leading to more lightning strikes, consequently damaging transmitters, and overhead cables, causing power outages. Increased cost of insurance for infrastructure in areas with repeated incidents of flooding, as well as withdrawal of risk coverage in vulnerable areas by private insurers. Road closures due to flooding thus inhibiting service and/or restoration efforts. Rising sea levels and corresponding increases in storm surges, increasing the risk of saline corrosion of coastal telecommunications infrastructure, and leading to erosion or inundation of coastal and underground infrastructure. 	<ul style="list-style-type: none"> Increased risk of storm surges impacting on coastal infrastructure. Increased storm intensity and frequency impacting on electricity and telecommunications infrastructure. 	(Naidoo, 2018)
Health	<ul style="list-style-type: none"> More exposure to high temperatures causing increased health risks including heat strokes. Heat waves increase threat of cardiovascular, kidney, and respiratory disorders. Increase in fire danger days causing increased loss of life and damage to health infrastructure. Wildfire smoke significantly reducing air quality, both locally and in areas downwind of fires. Smoke exposure increases respiratory and cardiovascular hospitalizations; emergency department visits; medication dispensations for asthma, bronchitis, chest pain, chronic obstructive pulmonary disease, and respiratory infections; and medical visits for lung illnesses. Increased emissions in biogenic volatile organic compounds from vegetation causing increases in air pollution. Increase in evaporative emissions from cars contributing to exposure to, and health impacts from, air pollution. Increase in distribution of vector-borne diseases in warmer areas. Increased water temperatures leading to an increase in algal blooms which can likely lead to increases in food- and waterborne exposures. Increased temperatures combined with fewer clouds (e.g., from increased subsidence that is projected for parts of South Africa) causing increased exposure to Information and Communication Technology which will have negative impacts on health. Increased temperatures increasing the reaction between certain pollutants and sunlight and heat, resulting in more severe hazardous smog events. 	<ul style="list-style-type: none"> Decreased soil moisture potentially creating more wind-blown dust which has negative impacts on air quality. Increase in water-borne diseases and diarrhoeal diseases due to inadequate water availability. Decreased precipitation causing changes in salinity of water, resulting in an increase in algal blooms which can likely lead to increases in food- and waterborne exposures. Increase in stagnant air, decreasing air quality. 	<ul style="list-style-type: none"> Wetter climate combined with increased temperatures may have negative health impacts as many diarrhoeal diseases vary seasonally, typically peaking during the rainy season. Extreme rainfall and higher temperatures increasing the prevalence of fungi and mould indoors, with increased associated health concerns. Increased flooding increasing the risk of drinking and wastewater treatment facilities being flooded, meaning that diarrhoeal diseases can be transmitted as wastewater systems overflow or drinking water treatment systems are breached. Increase in natural disasters (e.g., floods) creating a conducive environment for the occurrence of mental health problems. 	<ul style="list-style-type: none"> Increase in wind-blown dust combined with low humidity causing increased cases of meningitis (Davis, 2014). 	(Garland, 2018)

Parameter	Results of Climate Change				Reference
	Increase in temperature and heat stress	Drought and decrease in rainfall	Increase in rainfall and inland flooding	Increased wind speed	
Energy	<ul style="list-style-type: none"> Increased heat causing expansion of overhead cables, and cable sag. Sagging below a certain level result in a reduction in the amount of electricity transmitted. Increased heat stress on electricity transmission networks (overhead cables). Increase in heat island effect increasing energy demand for cooling, leading to grid stress. Increased threat of wildfires causing widespread damage to infrastructure and causing disruptions to service provision. 		<ul style="list-style-type: none"> Increase in flooding causing damage to electricity transmission and distribution infrastructure, poles, lines and sub-stations. Increase in frequency and cost of maintenance of concrete structures due to frequent and intense rainfall, flooding, or sea level rise. Increased repair events increasing stress put on service crews and resulting in delays to power restoration. 	<ul style="list-style-type: none"> Winds causing damage to energy supply infrastructure as winds cause overhead lines to sag, reducing electricity transmission. Extreme winds causing poles and trees to fall, causing further damage to energy supply infrastructure such as overhead lines. 	(Thambiran & van Wyk, 2018)
Ecosystem Services	<ul style="list-style-type: none"> Increased risks of water shortages increasing demand for irrigation of gardens and agriculture. Increased evapotranspiration rates with rising temperatures, reducing the water available in reservoirs and water available for reliant ecosystems. Increase in temperature leading to water loss via evapotranspiration resulting in decreased water quality and loss of wetlands. Loss or degradation of indigenous species, including threatened species or ecosystems. Increased threat from invasive species as competition for water increases. Dieback or death of susceptible plants (e.g., street trees) and animals (e.g., fish). Reduced availability of water and increased evapotranspiration resulting in reductions in harvested area (cropping area), yield (ton/ha) and quality. Warmer winters resulting in reduced period of dormancy (rest period) in deciduous fruit crops, decreasing the production and quality of associated food products. Warmer climate resulting in shifts in the growing season and life cycles of various plants, including crops, resulting in pests and diseases having a greater destructive impact as well as a shift in climatically suitable areas for specific crops. Increased humidity levels resulting in higher rates of microbial growth in fresh produce, reducing their expiry time. Increased heat stress on crops changes the micro-nutrients of crops products, decreasing the nutrient density and quality of food. Increased water temperature leading to increased growth of aquatic weeds which increases breeding of disease vectors and reduces water oxygen levels. Milder winters and reduced frost increase the duration of the growing season, increasing the survival rate of insects and diseases. Increased sea surface temperatures (SST) causing shifts in the spatial distribution of fish species. Increased SST and ocean acidification decreases marine phytoplankton growth and synthesis of omega-3 polyunsaturated fatty acids (PUFA's), affecting the oceanic food chain and consequent ecosystems. Increased heat stress and higher humidity levels 	<ul style="list-style-type: none"> Decreased amounts of rainfall reaching ecosystems as settlements use rainwater harvesting techniques for increased household use. Increased reliance on irrigation and greater demand for water to maintain public open space and gardens. Reduced planting and pollination leading to greater risk of erosion and soil loss. Increasing temperatures together with increased intensity of drought will potentially increase the occurrence of algal blooms in reservoirs and dams which are damaging to ecosystem functioning and water services. Drought and decreased rainfall causing wetland habitat loss. Locally specific changes in humidity levels will have impacts on local vegetation. Increased threat to watershed and aquifer recharge areas, affecting vegetation. Reduced soil moisture availability increasing moisture stress leading to dieback and death of plants and the loss or degradation of indigenous communities, including threatened species or ecosystems. Increased moisture stress leading to decline in crop yield and quality, and reduced fodder quantity and quality for livestock. Drying up of aquatic systems, perennial systems will become seasonal and seasonal systems will die off and be replaced by terrestrial plants. Increased spread of drought-adapted alien invasive plant species. 	<ul style="list-style-type: none"> Rainfall in shorter and more violent spells making recharging groundwater difficult. Increase in intensity of rainfall and flooding leading to increased surface runoff, resulting in increased soil erosion, soil loss and degradation. Increased rainfall and floods resulting in waterlogged soils which increase the likelihood of crop failure. Increasingly saturated soils leading to more standing water (ponding) which can result in more insect (pest) activity and their potential to carry diseases. Increased wave energy and run-up (sea level rise and more storms) causing degradation of natural coastal defence structures. 	<ul style="list-style-type: none"> Evapotranspiration rates increase with wind speed, reducing the water available in reservoirs and water available for reliant ecosystems. Increased rate of fire spread and spotting (the ignition of fires ahead of the main fire front) of fires. Potential damage to or uprooting of vegetation including trees, which can also damage infrastructure. Potential wind damage to crops, reducing yield and quality (e.g., sandblasting and fruit fall). Increased windblown materials (e.g., dust, litter) increasing the need for maintenance and city cleaning. Degradation of natural coastal defence structures and increased damage to hard coastal infrastructure. 	(Pieterse & Crankshaw, 2018)

Parameter	Results of Climate Change				Reference
	Increase in temperature and heat stress	Drought and decrease in rainfall	Increase in rainfall and inland flooding	Increased wind speed	
	potentially resulting in the exceedance of the temperature humidity index in livestock, causing reduced immunity, fertility, productivity and even mortality of livestock.				
Culture and Heritage	<ul style="list-style-type: none">• Increased temperature having significant impacts on the comfort levels of built heritage resources, resulting in the building no longer being fit-for-purpose.• Increased demand for additional heating and cooling resulting in the installation of heating, ventilation, and air-conditioning systems with potential negative consequences on the heritage value.• Increased heat stress potentially impacting on the materials and structural integrity of heritage resources.• Migration of several plant species due to changing climate patterns, posing a threat to the conservation of biodiversity hotspots, and potentially altering heritage places.• Increase in veld and forest fires raising the threat of fire to all heritage resources, natural and built, as well as posing health risks to heritage resource dwellers from exposure to smoke and ash pollution.	<ul style="list-style-type: none">• Decreased rainfall impacting negatively on ground moisture levels and thus the geological conditions of sensitive heritage resources. Drying out clays, for example, will shrink and potentially undermine founding conditions.	<ul style="list-style-type: none">• Increased rainfall in areas with clay soils resulting in swelling which poses a threat to the structural integrity of heritage resources.• Increased floods and changes in precipitation resulting in increasing vulnerability of archaeological evidence buried underground due to changing stratigraphic integrity of the soils.• Increased threat to materials and structural integrity of heritage resources exposed to higher humidity/precipitation levels.		(van Wyk, 2018)

4 IMPACT SIGNIFICANCE BASED ON SCOPING PHASE ASSESSMENT

The impact significance will be based on the quantified GHG emissions from the project. As the quantification of GHG for the project will only be undertaken during the impact assessment phase, the significance rating can only be provided once the impact assessment has been undertaken. No fatal flaws, however, are expected due to the climate impacts.

5 PLAN OF STUDY FOR THE IMPACT ASSESSMENT

During its construction, operation and closure phases the project will emit GHGs, the quantities of which will be estimated during the impact assessment phase of the project.

The impact assessment will include the following information:

- An estimation of the CO₂-equivalent (CO₂e) emissions from the project, associated fuel use, vegetation clearing activities, and electricity use;
- Estimate the impact of the project on national greenhouse gas emissions;
- Evaluation the potential impact of global climate change on the project by identifying potential physical risks to the project, employees, and communities;
- Provide the potential risk of climate change on the project and the risk of the project on climate change;
- Determination of environmental risk according to stipulated Impact Assessment methodology and,
- Recommendation of mitigation and management measures, where applicable.

The climate change impact assessment report will consider Scope 1 emissions, which are the emissions directly attributable to the proposed project, Scope 2 emissions, which are the emissions associated with bought-in electricity over the lifetime of the project, and Scope 3 emissions (as far as is reasonable and practically possible), which consider the “embedded” carbon in bought-in materials and downstream emissions.

6 REFERENCES

- Cole, M., Bailey, R., Cullis, J., & New, M. (2017). Spatial inequality in water access and water use in South Africa. *Water Policy*, 20 (1): 37–52.
- CSIR. (2025). Retrieved from Green Book: Adapting South African settlements to climate change.: www.greenbook.co.za
- Davis, C. E. (2014). *The International Traveler's Guide to Avoiding Infections*. Myers, J., Young, T., Galloway, M., Manyike, P. and Tucker, T. 2011 'References esponding to climate change in southern Africa – the role of research',.
- DFFE. (2022a). *GHG Inventory Report for South Africa 2000 -2020*. DFFE. Retrieved 09 14, 2022, from https://www.dffe.gov.za/sites/default/files/gazetted_notices/8thnational-greenhouse-gasinventory-reportg47133gon2321.pdf
- DFFE. (2022b). *Methodological Guidelines for Quantification of Greenhouse Gas Emissions: A companion to the South African National GHG Emission Reporting Regulations. Version No: MG-2022.1*. Pretoria: Government Gazette, 7 October 2022, No. 47257(2598), Department of Forestry, Fisheries and Environment.
- DFFE. (2025). *9th National Greenhouse Gas Inventory Report for the Republic of South Africa*. Pretoria, South Africa: Department of Forestry, Fisheries and Environment.
- Duncker, L. (2018). *Green Book – The impact of climate change on sanitation services provision within settlements*. Pretoria: CSIR.
- Dunker, L., & Van Wyk, L. (2018). *Green Book – The impact of climate change on stormwater management within settlements*. CSIR.
- Engelbrecht, F. (2019). *Technical Report: Green Book – Detailed Projections of Future. Report reference number: CSIR/BE/SPS/ER/2019/0005/C*. Pretoria: CSIR (Council for Scientific and Industrial Research).
- Garland, R. (2018). *Green Book – The impact of climate change on human health in settlements*. Pretoria: CSIR.
- Hofste, R., Kuzma, S., Walker, S., Sutanudjaja, E., Bierkens, M., Kuijper, M., . . . Reig, P. (2019). *Aqueduct 3.0: Updated Decision- Relevant Global Water Risk Indicators: Technical Note*. Washington, DC: World Resources Institute. Retrieved from Available online at: <https://www.wri.org/publication/aqueduct-30>
- IPCC. (2006). *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. Retrieved from Intergovernmental Panel on Climate Change: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/>
- IPCC. (2007). *Intergovernmental Panel on Climate Change 4th Assessment Report*. Retrieved from Intergovernmental Panel on Climate Change: http://www.ipcc.ch/pdf/assessment-report/ar4/syr/ar4_syr_appendix.pdf

- IPCC. (2013). *Intergovernmental Panel on Climate Change 5th Assessment Report*. Retrieved from Intergovernmental Panel on Climate Change: <https://www.ipcc.ch/report/ar5/>
- IPPC. (2022). *IPCC WGI Interactive Atlas: Regional information (Advanced)*. Retrieved January 6, 2023, from <https://interactive-atlas.ipcc.ch>
- Le Roux, A., van Niekerk, W., Arnold, K., Pieterse, A., Ludick, C., Forsyth, G., . . . Mans, G. (2019). *Green Book Risk Profile Tool*. (CSIR, Pretoria) Retrieved 09 30, 2022, from riskprofiles.greenbook.co.za
- Meteoblue. (2025, 08 14). *Climate Change*. Retrieved from meteoblue: <https://www.meteoblue.com/en/weather/historyclimate/change/-28.253N26.963E>
- Mokonyama, M., & Van Wyk, L. (2018). *Green Book - Transport and mobility: the impact of climate change on transport and mobility within settlements*. CSIR.
- Naidoo, S. (2018). *Green Book – The impact of climate change on ICT services provision within settlements*. . Pretoria: CSIR.
- NOAA. (2021, September). *Earth System Research Laboratory Global Monitoring Division*. Retrieved from National Oceanic and Atmospheric Administration: <https://gml.noaa.gov/ccgg/trends/global.html>
- Oelofse, S. (2018). *The impact of climate change on solid waste management within settlements*. CSIR.
- Pieterse, A., & Crankshaw, B. (2018). *Green Book –The impact of climate change on ecosystem services in settlements*. Pretoria: CSIR.
- SAWS. (2017). *A Climate Change Reference Atlas 2017 - based on CMIP5 – CORDEX downscaling*. Pretoria, South Africa: South African Weather Service and the Water Research Commission. Retrieved from <https://www.weathersa.co.za/home/climatechangeatlas>
- South African National Biodiversity Institute. (2025, 02 05). 2012 Vegetation Map App [Vector] 2012. Pretoria. Retrieved from Available from the Biodiversity GIS website: <http://bgis.sanbi.org/SpatialDataset/Detail/595>
- Thambiran, T., & van Wyk, L. (2018). *Green Book – The impact of climate change on energy services provision within settlements*. Pretoria: CSIR.
- UNFCCC. (2017). *United Nations Framework Convention on Climate Change e-Handbook* . Retrieved from United Nations Framework Convention on Climate Change: <http://bigpicture.unfccc.int/>
- van Wyk, L. (2018). *Green Book – The impact of climate change on culture and heritage in settlements*. Pretoria: CSIR.